arXiv:1612.06726v2 [math.AG] 21 Dec 2016

NODAL SURFACES WITH SINGULAR OR TOO LARGE DEFORMATION SPACES

REMKE KLOOSTERMAN

Abstract. In this text we show that the deformation space of a nodal surface S of degree d is smooth and of the expected dimension if $d \leq 7$ or $d \geq 8$ and S has at most $4d-5$ nodes. (The case $d \leq 7$ was previously covered by Alexandru Dimca, by different techniques.)

For $d \geq 8$ we give explicit examples of nodal surfaces with $4d-4$ nodes, for which the tangent space to the deformation space has larger dimension than expected.

1. Introduction

In his recent PhD thesis Yan Zhao [9] studied the deformation theory of nodal surfaces. He showed that for particular families of sextic surfaces the dimension of the deformation space equals the expected dimension. There are various deformation spaces for singular varieties (i.e., either preserving the analytic or the topological types of the singularities or the equisingular deformation space). However, for nodal varieties these spaces coincide and we call them "the deformation space".

Zhao's results motivated us to prove the following result.

Theorem 1.1. *Let $S \subset \mathbf{P}^3$ be a nodal surface of degree d. Suppose that $d \leq 7$ or $d \geq 8$ and S has at most $4(d-1)-1$ nodes then the deformation space of S is smooth and has the expected dimension.*

The above bound is sharp:

Theorem 1.2. *Suppose $S \subset \mathbf{P}^3$ is a nodal surface of degree $d \geq 8$ with $4(d-1)$ nodes. Then the tangent space to the deformation space does not have the expected dimension if and only if the locus of the nodes form a complete intersection of degree $(1, 4, d-1)$.*

The proof relies on Zhao's result that the difference between the expected dimension and the actual dimension of the tangent space at a surface S equals the defect of the linear system of degree d polynomials through the nodes of S. Then we use some ideas from [7, Proof of Theorem 4.1] to bound the Hilbert function of an ideal I, which has defect in certain degree, and has finite base locus in a lower degree.

As an application we discuss surfaces of the form

$$f_1 f_2 + f_3^2 f_4$$

with $\deg(f_1) = 1$ and $4 \leq \deg(f_3) \leq d/2$. (See Example 2.6.)

The author would like to thank Arnaud Beauville for pointing out the paper [3].

Deformation theory of singular hypersurfaces have been studied extensively. The starting point for deformation of nodal surfaces is the paper [1]. For our applications, the slightly different presentation in [9, Chapter 3] is more suitable.

2. Proof of the theorems

Let $S = \mathbf{C}[x_0, \ldots, x_3]$. For a homogeneous ideal $I \subset S$ denote with $h_I(k) = \dim S/I_k$, the Hilbert function of I.

Macaulay [8] described the possible Hilbert functions of homogeneous ideals. Gotzmann [5] described what happens in the extreme case. As a corollary we obtain the following result. (For the deduction see [7, Section 2].)

Theorem 2.1. *Let $I \subset S$ be an ideal. If $h_I(k) \leq k$ then $h_I(k+1) \leq h_I(k)$. If, moreover, I_{k+1} is base point free then $h_I(k+1) < h_I(k)$ or $h_I(k) = 0$.*

This result is a key ingredient in various proofs of the explicit Noether-Lefschetz Theorem (e.g. [6]) and is also used in the author's proof [7] for the fact that a nodal hypersurfaces of degree d in $\mathbf{P}^4$ with less than $(d-1)^2$ nodes is factorial.

In the following we use the Alexander polynomial of hypersurfaces with isolated singularities. Suppose $X = V(f) \subset \mathbf{P}^n$ is a hypersurface with isolated singularities. Then f defines an affine hypersurface with one-dimensional singularities. Let $F = V(f-1) \subset \mathbf{A}^{n+1}$ be the affine Milnor fiber. Fix a primitive d-th root of unity ζ. Let T be the map multiplying each x_i with ζ. Then T acts on F. The Alexanderpolynomial Δ_X of X is the characteristic polynomial of T^* acting on $H^{n-1}(F)$. The only possible zeros are d-th roots of unity. One can bound the exponent of $(t-\alpha)$ by $(-1)^{n+1}\frac{1-(-1)^n(d-1)^n}{d}$, if α is a d-th root of unity different from one. In the case $n = 3$ we find that the exponent is at most $d^2 - 3d + 1$. For more on this see [2].

The following proposition is the same as [3, Corollary 1.3], but out proof is different from Dimca's proof.

Proposition 2.2. *Let S be a nodal surface of degree d. Let Σ be the locus of points where S has a node and let $I = I(\Sigma)$. Then $h_I(k) = \#\Sigma$ for $k \geq \frac{3}{2}d - 3$.*

In particular, if $h_I(d) < \#\Sigma$ then $d \geq 8$.

Proof. Recall that the Alexander polynomial of a nodal surface equals $(t+1)^\delta$ where δ is the defect of the linear system of degree $\frac{3}{2}d - 4$ polynomials through Σ. (E.g., [2, Theorem 6.4.5])

Let

$$0 \to \oplus_{j=0}^{e_k} S(-b_{k,j}) \to \oplus_{j=0}^{e_{k-1}} S(-b_{k-1,j}) \to \cdots \to \oplus_{j=0}^{e_0} S(-b_{0,j}) \to S/I \to 0$$

be a minimal resolution of S/I. Let $\beta_m := \sum_{j=0}^{k} (-1)^j \#\{i | b_{j,i} = m\}$. From [4, Theorem 4.2] it follows that $\beta_m \neq 0$ for $m = \max b_{j,i}$. An easy calculation shows that the defect δ equals

$$\sum_{m \geq \frac{3}{2}d} \beta_m \binom{m - \frac{3}{2}d + 3}{3}$$

if d is even.

Let s be an even integer. Take a general degree-s base change $\varphi : \mathbf{P}^3 \to \mathbf{P}^3$. Let $\tilde{I}$ be the ideal of the nodes of $\varphi^{-1}(S)$.The pull back of the resolution of S/I is a minimal resolution of $S/\tilde{I}$.

In particular, we see that δ associated with $\varphi^{-1}(S)$ equals

$$\sum_{m\geq\frac{3}{2}d}\beta_m\binom{s(m-\frac{3}{2}d)+3}{3}.$$

If $\beta_m\neq 0$ for some $m>\frac{3}{2}d$ then δ (as a function in s) behaves asymptotically as s^3 times a constant. However, as we argued above the exponents occuring in the Alexander polynomial can be bounded by a function of type $(sd)^2$ times a constant. Hence $\beta_m=0$ for all $m>\frac{3}{2}d$ and therefore $h_I(k)=\#\Sigma$ for $k\geq\frac{3}{2}d-3$ and, if d is even, $h_I(\frac{3}{2}d-4)=\#\Sigma-\beta_{3d/2}$. □

Remark 2.3. This proposition is the same as [3, Corollary 1.3]. Dimca bounds the degree of the syzygies by using spectral sequences and an argument comparing the Hodge filtration and the pole order filtration on the cohomology of the complement of the affine Milnor fiber associated with the defining polynomial of S.

Proposition 2.4. *Let S be a nodal surface of degree d. Suppose $d\geq 8$. Let I be the ideal of the nodes of S. If S has at most $4d-5$ nodes then $h_I(d)=\#\Sigma$. If S has $4d-4$ nodes and $h_I(d)<\#\Sigma$ then I is a complete intersection ideal of degree $(1,4,d-1)$.*

Proof. Let ℓ be a general linear form. Let $I_H=(I,\ell)$. Consider the exact sequence

$$0\to(S/I)_k\xrightarrow{\ell}(S/I)_{k+1}\to(S/I_H)_{k+1}\to 0.$$

Denote with $I_H=(I,\ell)$. From this exact sequence it follows that

$$h_I(k)=\sum_{j=0}^{k}h_{I_H}(k).$$

The vector space I_{d-1} contains all partial derivates of the defining polynomial of S. In particular it has a finite base locus and therefore $(I_H)_{d-1}$ has empty base locus. Suppose now that $h_I(d)<\#\Sigma$. Since $h_I(k)=\#\Sigma$ for k large we get that $h_{I_H}(k)\neq 0$ for some $k>d$ and therefore that $h_{I_H}(k)\neq 0$ for $0\leq k\leq d+1$. Let $J\subset S$ be the following ideal. Take J_{d+1} such that $(I_H)_{d+1}\subset J_{d+1}$ and $h_J(d+1)=1$. For $k<d+1$ set

$$J_k=\{f\in S_k\colon fS_{d+1-k}\subset J_{d+1}\}$$

and $J_k=S_k$ for $k>d+1$. Then J is an ideal and S/J is Artinian Gorenstein with socle degree $d+1$. In particular, $h_J(k)=h_J(d+1-k)$.

Since J_{d-1} is base point free and $d-2>4$ we can use Theorem 2.1 to obtain $h_J(d)\geq 2, h_J(d-1)\geq 3$ and $h_J(d-2)\geq 4$. Similarly, we find that $h_J(k)\geq 4$ for $k\geq 4$. Using the duality we get that $h_J(0)=1, h_J(1)\geq 2, h_J(2)\geq 3$ and $h_J(3)\geq 4$. All in all we find

$$\#\Sigma\geq h_I(d+1)=\sum_{k=0}^{d+1}h_{I_H}(k)\geq\sum_{k=0}^{d+1}h_J(k)\geq 4d-4.$$

Hence if $\#\Sigma<4d-4$ then $h_I(d)=\#\Sigma$.

If $\#\Sigma<h_I(d)$ and $\#\Sigma=4(d-1)$ hold then we have equality everywhere, i.e., $J=I_H$. We also have complete knowledge of $h_I(k)$ for $k\leq d$ and we find that I has a generator in degree 1 and in degree 4. Moreover we have that $h_I(k)$ equals the Hilbert function of a plane curve of degree 4, for $k<d-1$. Hence there are no

further generators in degrees up to $d-2$. However, we have one further generator in degree $d-1$. Since I_{d-1} has finite base locus, we have that these three generators form a complete intersection. The length of this complete intersection is $1\cdot 4\cdot(d-1)$. Hence Σ is a complete intersection of degree $(1,4,d-1)$. □

Remark 2.5. In the case that $h_I(d) < \#\Sigma$ and $\#\Sigma = 4(d-1)$ we have that $I = (g_1, g_2, g_3)$, with $\deg(g_1) = 1, \deg(g_2) = 4, \deg(g_3) = d-1$. Since I is a complete intersection ideal we have that I^2 is generated by products of the generators of I. From $f \in I^2$ and $\deg(g_3^2) > \deg(f)$ it now follows that $f \in (g_1, g_2)$, i.e., the quartic curve containing all the nodes is contained in S.

Proof of the Theorems. Let $S \subset \mathbf{P}^3$ be a nodal surface. In [9, Chapter 3] shows that the tangent space to the deformation space of S can be identified with $(I/J)_d$, where I is the ideal of the nodes of S and J is the Jacobian ideal of S. Then [9, Proposition 3.3.8] implies that $\dim J_d = 16$. In particular, the dimension of the tangent space to the deformation space of S has the expected dimension if and only if the ideal of the nodes of S does not have defect in degree d.

Hence if $d < 8$ or $d \geq 8$ and $\#\Sigma < 4d-4$ then the tangent space has the expected dimension and therefore the deformation space has the expected dimension and is smooth. If $d \geq 8$ and $\#\Sigma = 4d-4$ holds and the tangent space has too large dimension if and only if the ideal of the nodes is a complete intersection of degree $(1,4,d-1)$. □

Example 2.6. Let $d \geq 8$ and let $4 \leq a \leq \frac{1}{2}d$.

Fix general polynomials f_1, f_2, f_3, f_4 of degree $1, d-1, a, d-2a$. Then

$$f_1 f_2 + f_3^2 f_4$$

defines a hypersurface with singularities at $f_1 = f_2 = f_3 = 0$. If the forms are chosen general then there are no further singularities and all singular points are nodes. Hence we a have that the locus of the nodes is a complete intersection of degree $(1, a, d-1)$. We have that $h_I(k) < a(d-1)$ if and only if $k \leq d+a-4$. In particular $h_I(d) < \#\Sigma$.

The locus L of polynomials of this shape has dimension

$$\binom{d-1+3}{3} + 4 - 1 + \binom{a+2}{2} + \binom{d-2a+2}{2} - 1$$

This implies that the codimension of L in S_d equals

$$\frac{1}{2}(-5a^2 + 3a + 4da - 6).$$

For small a this codimension is lagger then the number of nodes. Hence for small a we have that L is a proper closed subset of some component of the deformation space.

For

$$\frac{1}{10}2d + 5 + \sqrt{4d^2 + 20d - 95} < a < \frac{1}{2}d$$

we have that the codimension of L in S_d is less then $a(d-1)$. However, the codimension of L in S_d is strictly larger than $h_I(d)$. So either L is a component of the deformation space, and this component is nonreduced or L is a proper closed subset in a larger component.

Finally, if d is even and $a = \frac{1}{2}d$ then the codimension of L in S_d equals $h_I(d)$. In particular, L is an irreducible component of the deformation space and this component is smooth and *not* of the expected dimension.

We have two geometric-topological characterisations for the case $a = \frac{1}{2}d$.

If $a < 1/2d$ then we have that $\operatorname{rank} \mathrm{CH}^1(S) > 1$, (i.e., the hyperplane section and the degree a curve $f_1 = f_3 = 0$ are linearly independent in $\mathrm{CH}^1(S)$), whereas for $a = \frac{1}{2}d$ we have $\operatorname{rank} \mathrm{CH}^1(S) = 1$ for very general S.

A second characterisation is given by the Alexander polynomial, this polynomial is a constant polynomial in the case $a < 1/2$ and equals $(t+1)$ in the case $a = \frac{1}{2}d$.

References

[1] D. M. Burns, Jr. and J. M. Wahl. Local contributions to global deformations of surfaces. *Invent. Math.*, 26:67–88, 1974.

[2] A. Dimca. *Singularities and topology of hypersurfaces*. Universitext. Springer-Verlag, New York, 1992.

[3] A. Dimca. On the syzygies and Hodge theory of nodal hypersurfaces. Preprint available at `arxiv:math/1310.5344v1`, 2013.

[4] D. Eisenbud. *The geometry of syzygies*, volume 229 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.

[5] G. Gotzmann. Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes. *Math. Z.*, 158:61–70, 1978.

[6] M.L. Green. Components of maximal dimension in the Noether-Lefschetz locus. *J. Differential Geom.*, 29:295–302, 1989.

[7] R. Kloosterman. Maximal families of nodal varieties with defect. Preprint, available at `arXiv:1310.0227v1`, 2013.

[8] F. S. Macaulay. Some Properties of Enumeration in the Theory of Modular Systems. *Proc. London Math. Soc.*, S2-26:531, 1927.

[9] Y. Zhao. *Deformations of nodal surfaces*. PhD thesis, Leiden University, 2016. Available at `https://openaccess.leidenuniv.nl/handle/1887/44549`.

Università degli Studi di Padova, Dipartimento di Matematica, Via Trieste 63, 35121 Padova, Italy